Efficacy of deucravacitinib, an oral, selective, allosteric tyrosine kinase 2 (TYK2) inhibitor, in scalp psoriasis by baseline Psoriasis Area and Severity Index (PASI) and baseline body surface area (BSA): a subanalysis of the phase 3 clinical trial data


Andrew Blauvelt, Howard Sofen, Jo Lambert, Joseph F. Merola, Mark Lebwohl, Kim Hoyt, Renata M. Kisa, Subhashis Banerjee, Thomas Scharnitz, Jeffrey J. Crowley

¹Oregon Medical Research Center, Portland, OR, USA; ²University of California Los Angeles and Dermatology, Bakersfield Dermatology, Bakersfield, CA, USA; ¹Ghent University, Ghent, Belgium; ⁴UT Southwestern Medical Center, Dallas, TX, USA; ¹Bristol Myers Squibb, Princeton, NJ, USA; ¹Bristol Myers Squibb, Princeton, NJ, USA; ¹Bakersfield Dermatology, Bakersfield, CA, USA

Introduction

- Tyrosine kinase 2 (TYK2) is an intracellular enzyme that mediates signaling of select inflammatory cytokines (eg, interleukin [IL]-23, IL-12, Type I interferons [IFNs])
- IL-23 and Type I IFNs are involved in psoriasis pathogenesis
- Deucravacitinib, an oral, selective, allosteric TYK2 inhibitor, is approved in the US, EU, and other countries for the treatment of adults with moderate to severe plaque psoriasis who are candidates for systemic therapy^{2.6}
- Deucravacitinib uniquely binds to the TYK2 regulatory domain rather than to the catalytic domain where Janus kinase (JAK) 1,2,3 inhibitors bind^{1,7} (Figure 1), driving its selectivity for TYK2 and representing the first in a new class of oral drugs

Figure 1. Mechanism of action of deucravacitinib



- Deucravacitinib demonstrated a robust efficacy and safety profile in the global phase 3 POETYK PSO-1 (NCT03624127) and POETYK PSO-2 (NCT03611751) trials in patients with moderate to severe plague psoriasis
- Deucravacitinib was superior to placebo and apremilast at Week 16, maintained response rates through Week 52, and was well tolerated8,
- Deucravacitinib demonstrated efficacy in scalp psoriasis:
- In patients with moderate to severe scalp psoriasis (scalp-specific Physician Global Assessment [ss-PGA] score ≥3) at baseline in POETYK PSO-1 and PSO-2, deucravacitinib was superior to placebo based on scalp-specific endpoints (ss-PGA score of 0 [clear] or 1 [almost clear] [ss-PGA 0/1], ≥90% reduction from baseline in Psoriasis Scalp Severity Index [PSSI 90])11
- $-\,$ ss-PGA 0/1 response rates were maintained through Week 52 in patients who continued to receive deucravacitinib1
- · Efficacy in scalp psoriasis was observed regardless of body weight, body mass index, or type or number of prior psoriasis therapies
- However, the effect of baseline disease severity overall on deucravacitinib efficacy in scalp psoriasis has

· To evaluate the effect of baseline disease severity overall on the efficacy of deucravacitinib in scalp psoriasis in the phase 3 POETYK PSO-1 and PSO-2 trials

Methods

POETYK PSO-1 and PSO-2 study designs

- POETYK PSO-1 and PSO-2 were global, 52-week, phase 3, double-blind trials that randomized adults with moderate to severe plaque psoriasis 1:2:1 to oral placebo, deucravacitinib 6 mg once daily (QD), or apremilast 30 mg twice daily8,9
- Placebo patients crossed over to deucravacitinib at Week 16
- In POETYK PSO-2, deucravacitinib-treated patients who achieved ≥75% reduction from baseline in Psoriasis Area and Severity Index (PASI 75) at Week 24 were rerandomized 1:1 to continue deucravacitinib

Analysis populations

- Patients with moderate to severe scalp psoriasis at baseline (ss-PGA ≥3) were evaluated in the following
- Pooled POETYK PSO-1/PSO-2 patients who were treated with deucravacitinib 6 mg QD through Week 24 or who crossed over from placebo to deucravacitinib at Week 16
- POETYK PSO-1 patients who received continuous deucravacitinib 6 mg QD without change from Day 1 through Week 52 and patients who crossed over from placebo to deucravacitinib at Week 16
- Patients in the 2 populations were stratified by baseline disease severity:
- Baseline body surface area (BSA) involvement: 10%-≤15% and >15% Baseline PASI score: 12-<15 and ≥15
- Scalp-specific outcomes in the pooled POETYK PSO-1/PSO-2 population through Week 24:
- ss-PGA 0/1 PSSI 90
- Scalp-specific outcomes in the POETYK PSO-1 population through Week 52:
- PSSI 90

- Nonresponder imputation (NRI) was used for binary endpoints for patients who discontinued early or had
- Confidence intervals (CIs) were obtained using a stratified Cochran-Mantel-Haenszel test for pooled POETYK PSO-1/PSO-2 with stratification factors for study and using the Clopper-Pearson method for POETYK PSO-1

Results

Baseline patient demographics

- · Baseline patient demographics were generally similar across the various disease severity subgroups in the pooled POETYK PSO-1/PSO-2 (Table 1) and POETYK PSO-1 (Table 2) populations
- Table 1. Baseline patient demographics (pooled POETYK PSO-1 and PSO-2)

Parameter	Baseline BSA involvement				Baseline PASI score			
	Placebo		Deucravacitinib		Placebo		Deucravacitinib	
	10%-≤15% (n = 95)	>15% (n = 199)	10%-≤15% (n = 147)	>15% (n = 367)	12-<15 (n = 76)	≥15 (n = 218)	12-<15 (n = 96)	≥15 (n = 418)
Age, mean (SD), y	47.3 (14.7)	46.3 (14.3)	45.8 (14.2)	44.0 (13.4)	46.4 (15.5)	46.7 (14.0)	45.5 (13.4)	44.5 (13.7)
Weight, mean (SD), kg	88.5 (21.0)	90.9 (21.6)	89.4 (21.8)	91.6 (22.5)	90.9 (20.7)	89.9 (21.6)	84.5 (20.6)	92.5 (22.4
Body mass index, mean (SD), kg/m²	29.8 (6.6)	30.4 (6.9)	29.7 (6.2)	31.0 (7.3)	30.2 (6.7)	30.2 (6.9)	29.0 (7.8)	31.0 (6.8)
Female, n (%)	36 (37.9)	63 (31.7)	51 (34.7)	131 (35.7)	25 (32.9)	74 (33.9)	37 (38.5)	147 (34.7)
Race, n (%)								
White	80 (84.2)	172 (86.4)	133 (90.5)	305 (83.1)	67 (88.2)	185 (84.9)	82 (85.4)	356 (85.2)
Asian	9 (9.5)	21 (10.6)	10 (6.8)	51 (13.9)	6 (7.9)	24 (11.0)	13 (13.5)	48 (11.5)
Other	6 (6.3)	6 (3.0)	4 (2.7)	11 (3.0)	3 (3.9)	9 (4.1)	1 (1.0)	14 (3.4)

Table 2. Baseline patient demographics (POETYK PSO-1)

		Baseline BSA	involvement		Baseline PASI score				
	Deucravacitinib		Placebo to deucravacitinib		Deucravacitinib		Placebo to deucravacitinib		
Parameter	10%-≤15% (n = 47)	>15% (n = 145)	10%-≤15% (n = 36)	>15% (n = 70)	12-<15 (n = 30)	≥15 (n = 162)	12-<15 (n = 29)	≥15 (n = 77)	
Age, mean (SD), y	46.6 (14.9)	43.7 (13.8)	46.3 (14.4)	46.8 (14.9)	42.4 (13.9)	44.7 (14.1)	44.8 (15.0)	47.3 (14.6)	
Weight, mean (SD), kg	91.7 (23.3)	87.7 (23.7)	88.3 (20.3)	88.4 (22.5)	85.9 (22.0)	89.2 (23.9)	91.1 (21.7)	87.3 (21.8)	
Body mass index, mean (SD), kg/m²	30.2 (6.4)	30.0 (8.1)	30.2 (6.1)	29.8 (7.1)	30.0 (9.7)	30.1 (7.3)	30.3 (6.8)	29.8 (6.8)	
Female, n (%)	14 (29.8)	50 (34.5)	12 (33.3)	24 (34.3)	12 (40.0)	52 (32.1)	9 (31.0)	27 (35.1)	
Race, n (%)									
White	41 (87.2)	107 (73.8)	28 (77.8)	53 (75.7)	22 (73.3)	126 (77.8)	24 (82.8)	57 (74.0)	
Asian	5 (10.6)	35 (24.1)	6 (16.7)	16 (22.9)	8 (26.7)	32 (19.8)	4 (13.8)	18 (23.4)	
Other	1 (2.1)	3 (2.1)	2 (5.6)	1 (1.4)	0	4 (2.5)	1 (3.4)	2 (2.6)	

dy surface area: PASI, Psoriasis Area and Severity Index: SD, standard deviation

- In the pooled POETYK PSO-1/PSO-2 population, ss-PGA 0/1 response rates were higher with deucravacitinib than with placebo at Week 16 in both baseline BSA involvement subgroups (Figure 2)
- ss-PGA 0/1 response rates were maintained with deucravacitinib in both subgroups through Week 24
- In POFTYK PSO-1. ss-PGA 0/1 response rates were maintained in both BSA involvement subgroups through Week 52 among patients who received continuous deucravacitinib treatment from Day 1 (Figure 3)
- Patients who crossed over from placebo to deucravacitinib at Week 16 achieved similar ss-PGA 0/1
- response rates at Week 52 compared with those receiving continuous deucravacitinib treatmen

Figure 2. ss-PGA 0/1 response rates through Week 24 by baseline BSA involvement (pooled POETYK PSO-1/PSO-2, NRI)





Figure 3. ss-PGA 0/1 response rates through Week 52 by baseline BSA involvement



ss-PGA 0/1 by baseline PASI score

- In the pooled POETYK PSO-1/PSO-2 population, ss-PGA 0/1 response rates were higher with deucravacitinib than with placebo at Week 16 in both baseline PASI score subgroups (Figure 4)
- ss-PGA 0/1 response rates were maintained with deucravacitinib in both subgroups through Week 24
- In POETYK PSO-1, ss-PGA 0/1 response rates were maintained in both PASI score subgroups through Week 52
- among patients who received continuous deucravacitinib treatment from Day 1 (Figure 5)
- Patients who crossed over from placebo to deucravacitinib at Week 16 achieved similar ss-PGA 0/1 response rates at Week 52 compared with those receiving continuous deucravacitinib treatmen

Figure 4. ss-PGA 0/1 response rates through Week 24 by baseline PASI score (pooled POETYK PSO-1/PSO-2, NRI)





Figure 5. ss-PGA 0/1 response rates through Week 52 by baseline PASI score (POETYK PSO-1, NRI)



PSSI 90 by baseline BSA involvement

- In the pooled POETYK PSO-1/PSO-2 population, PSSI 90 response rates were higher with deucravacitinib than with placebo at Week 16 in both baseline BSA involvement subgroups (Figure 6)
- PSSI 90 response rates were maintained with deucravacitinib in both subgroups through Week 24 • In POETYK PSO-1, PSSI 90 response rates were maintained in both BSA subgroups through Week 52 among patients who received continuous deucravacitinib treatment from Day 1 (Figure 7)
- Patients who crossed over from placebo to deucravacitinib at Week 16 achieved similar PSSI 90 response rates at Week 52 compared with those receiving continuous deucravacitinib treatment

Figure 6. PSSI 90 response rates through Week 24 by baseline BSA involvement (pooled POETYK PSO-1/PSO-2, NRI)





Figure 7. PSSI 90 response rates through Week 52 by baseline BSA involvement



PSSI 90 by baseline PASI score

- In the pooled POETYK PSO-1/PSO-2 population, PSSI 90 response rates were higher with deucravacitinib than with placebo at Week 16 in both baseline PASI score subgroups (Figure 8)
- PSSI 90 response rates were maintained with deucravacitinib in both subgroups through Week 24
- In POETYK PSO-1, PSSI 90 response rates were maintained in both PASI subgroups through Week 52 among patients who received continuous deucravacitinib treatment from Day 1 (Figure 9
- Patients who crossed over from placebo to deucravacitinib at Week 16 achieved similar PSSI 90 response rates at Week 52 compared with those receiving continuous deucravacitinib treatmer

Figure 8. PSSI 90 response rates through Week 24 by baseline PASI score (pooled POETYK PSO-1/PSO-2, NRI)





Figure 9. PSSI 90 response rates through Week 52 by baseline PASI score (POETYK PSO-1, NRI



Conclusions

- In POETYK PSO-1 and PSO-2, deucravacitinib improved scalp-specific disease burden regardless of overall baseline disease severity (baseline BSA involvement and PASI score) in patients with moderate
 - Higher ss-PGA 0/1 and PSSI 90 response rates were observed at Week 16 in patients receiving deucravacitinib compared with placebo
- Response rates were maintained through Week 52 in patients receiving continuous deucravacitinib treatment from baseline and were increased through Week 52 in patients crossing over from placebo to deucravacitinib at Week 16
- Patients with higher disease burden (BSA >15%; PASI ≥15) performed numerically better than those with lower disease burden; however, both groups demonstrated improvement relative to placebo and had continued improvement over time
- These findings suggest that deucravacitinib treatment improves scalp-specific disease regardless of baseline disease severity overall in patients with moderate to severe plague psoriasis
- Results are consistent with the previous report from POETYK PSO-1 and PSO-2 in which deucravacitinil also improves scalp psoriasis regardless of body weight, body mass index, or use of prior psoriasis

References

- 1. Burke JR, et al. Sci Transl Med. 2019;11:eaaw1736.
- 2. Sotyktu [package insert]. Princeton, NJ, USA: Bristol Myers Squibb; September 2022.
- 3. Sotyktu [summary of product characteristics]. Dublin, Ireland: Bristol Myers Squibb Pharmaceutical Operations; March 2023
- 4. Sotyktu [package insert]. Tokyo, Japan: Bristol Myers Squibb K.K.; September 2022
- 5. Sotyktu [product information]. Mulgrave, VIC, Australia: Bristol Myers Squibb Australia Pty. Ltd.; December 2022.
- 6. Sotyktu [product monograph], Montreal, OC, Canada: Bristol Myers Squibb Canada Co.: November 2022.
- 7. Wrobleski ST, et al. J Med Chem. 2019:62:8973-8995.
- 8. Armstrong AW, et al. J Am Acad Dermatol. 2023;88:29-39.
- 9. Strober B. et al. J Am Acad Dermatol. 2023:88:40-51.
- 10. Blauvelt A, et al. J Am Acad Dermatol, 2023 Dec 18:S0190-9622(23)03374-1, doi:10.1016/j.iaad.2023.11.060, Online ahead

Acknowledgments

- · This study was sponsored by Bristol Myers Squibb
- Writing and editorial assistance was provided by Jieming Fang, MD, of Peloton Advantage, LLC, an OPEN Health company,

 AB: Speaker (with honoraria): AbbVie, Bristol Myers Squibb, Lilly, Pfizer, Regeneron, and Sanofi: Scientific adviser (with honoraria): AbbVie, Abcentra, Aclaris, Affibody, Aligos, Almirall, Alumis, Amgen, AnaptysBio, Apogee, Arcutis, Arena, Aslan, Athenex, Bluefin Biomedicine, Boehringer Ingelheim, Bristol Myers Squibb, Cara Therapeutics, CTI BioPharma, Dermavant, EcoR1, Escient, Evelo Biosciences, Evommune, Forte Biosciences, Galderma, Highlightll Pharma, Incyte, InnoventBio, Jansser Landos, Leo Pharma, Lilly, Lipidio, Merck, Monte Rosa Therapeutics, Nektar, Novartis, Overtone Therapeutics, Paragon, Pfizer Rani, Rapt, Regeneron, Sanofi Genzyme, Spherix Global Insights, Sun Pharma, Takeda, TLL Pharmaceutical, TrialSpark, UCB,

AbbVie, Acelyrin, Allakos, Almirall, Alumis, Amgen, Arcutis, Athenex, Boehringer Ingelheim, Bristol Myers Squibb, Concert, Sanofi, Sun Pharma, UCB, and Ventyx Biosciences

Union, Ventyx Biosciences, Vibliome, and Xencor; Clinical study investigator (institution has received clinical study funds):

JL: Unrestricted grants: AbbVie, Almirall, Amgen/Celgene, Janssen-Cilag, Leo Pharma, Lilly, Novartis, and UCB; Speaker: AbbVie, Almirall, Bristol Myers Squibb, Janssen-Cilag, Pfizer, and UCB; Consultant: AbbVie, Amgen/Celgene, Bristol Myers

• JFM: Consultant and/or investigator: AbbVie, Amgen, Biogen, Bristol Myers Squibb, Dermavant, Janssen, Leo Pharma, Lilly, Novartis, Pfizer, Regeneron, Sanofi, Sun Pharma, and UCB

• ML: Research funds on behalf of Mount Sinai: AbbVie, Amgen, Arcutis, Avotres, Boehringer Ingelheim, Cara Therapeutics, Dermavant, Incyte, Janssen, Lilly, Ortho Dermatologics, Regeneron, and UCB; Consultant: Almirall, AltruBio, AnaptysBio Arcutis. Avotres. Boehringer Ingelheim. Brickell Biotech, Bristol Myers Squibb, Castle Biosciences, Celltrion, CorEvitas Psoriasis legistry, Dermavant, EPI Health, Evommune, Forte Biosciences, Galderma, Genentech, Incyte, Leo Pharma, Meiji Seika Pharma, Mindera Health, Pfizer, Seanergy, Strata, Trevi, and Verrica

- KH: Consultant: Bristol Myers Squibb via Syneos Health
- RMK, SB, and TS: Employees and shareholders: Bristol Myers Squibb
- JJC: Consultant, speaker, and/or investigator: AbbVie, Arcutis, Boehringer Ingelheim, Bristol Myers Squibb, Janssen, Lilly, Regeneron, Sanofi, and UCB